

Head and Neck Cancers

Gabriele Price
Oxford Cancer Intelligence Unit
18 May 2011




√ The Head and Neck Cancers e-Atlas

- ✓ Profile of Head and Neck Cancers in England: Incidence, Mortality and Survival
- ✓ Oral Cavity Cancer: Survival Trends in England






√ The Head and Neck Cancers e-Atlas




- ➤ Incidence and mortality
- ➤8 head and neck cancer subtypes: oral cavity, salivary glands, oropharynx, nasopharynx, hypopharynx, larynx, palate, thyroid gland





- ✓ Trends in incidence and mortality in England between 1990 to 2006
- ✓ Incidence and mortality by CN and SHA, 2002-06
- √ 1-year and 5-year relative survival in England, 1990-92 to 2004-06







- The incidence of several head and neck cancers has risen between 1990 and 2006:
 - o oral cavity
 - o salivary gland
 - o palate
 - o thyroid
 - o oropharyngeal












- The incidence of several head and neck cancers has risen between 1990 and 2006
- ➤ Mortality from several types of head and neck cancer types has fallen between 1990 and 2009:
 - nasopharyngeal
 - hypopharyngeal
 - laryngeal


Trends in the mortality of hypopharynx cancer in England, 1990–2006



















- The incidence of several head and neck cancers has risen between 1990 and 2006
- Mortality from several types of head and neck cancer types has fallen between 1990 and 2009
- ➤ Several cancer types have shown an improvement in relative survival rates during the study period:
 - oral cavity
 - salivary gland
 - o oropharyngeal
 - nasopharyngeal
 - o thyroid gland
 - hypopharyngeal
















Oral Cavity Cancer

Read NCIN's data briefing on Oral Cavity Cancer: Survival Trends in England




✓ Survival rates in men and women diagnosed in the period 2004 to 2006 across areas with varied deprivation (IMD 2007) levels

Trends in 1-year and 3-year relative survival for men and women in England diagnosed with oral cavity cancer between 1990-92 and 2005-07









0%

Most deprived

■1-year (women)

3-year (women)



Most deprived

0%



Least deprived

■1-year(men) ■3-year(men)

Least deprived

Any questions?

Gabriele.Price@sph.nhs.uk www.ociu.nhs.uk www.ncin.org.uk

